


# Hunting Animalcula with Flavour in the LHC Era



*Andrzej J. Buras*  
*(Technical University Munich, TUM-IAS)*



**“Present and Future of B Physics”**  
**Cracow, January 2012**



# Overture

# 1676

A very important year for  
the humanity !

# 1676 : The Discovery of the Microuniverse (Animalcula) (The Empire of Bacteria)



**Antoni van Leeuwenhoek**  
**\*24.10.1632 †27.08.1723**

$10^{-6}\text{m}$

**~500 Microscopes**

(Magnification  
by ~300)


# Animalcula Hunters




**Antoni van Leeuwenhoek**  
**\*24.10.1632 †27.08.1723**



**Lazzaro Spallanzani**  
**\*12.01.1729 †12.02.1799**



**L. Pasteur**  
**\*27.12.1822 †28.09.1895**



**Robert Koch**  
**\*11.12.1843 †27.05.1910**

# An Excursion towards the Very Short Distance Scales:

1676 - 2020

Microuniverse

$10^{-6}\text{m}$

Bacteriology  
Microbiology

Nanouniverse

$10^{-9}\text{m}$

Nanoscience

Femtouniverse

$10^{-15}\text{m}$

Nuclear Physics  
Low Energy Elementary  
Particle Physics

Attouniverse

$10^{-18}\text{m}$

High Energy Particle  
Physics (present)

High Energy Proton-Proton  
Collisions at the LHC

$5 \cdot 10^{-20}\text{m}$

Frontiers of Elementary  
Particle Physics in 2010's

High Precision Measurements  
of Rare Processes (Europe,  
Japan, USA)

$10^{-21}\text{m}$

Zeptouniverse

# **Most important Message from this Talk**

**Antoni van Leeuwenhook discovered in 1676**

**Animalcula**

# Most important Message from this Talk

Antoni van Leeuwenhook discovered in 1676

**Animalcula**

We all expect to discover **New Animalcula**

in the coming years with the help

of **LHC**

and

**High Precision Experiments**

**But how will these  
New Animalcula look like ?**

**But how will these  
New Animalcula look like ?**

**Overture Completed!**

# Cracow Symphony No. 7

# **Cracow Symphony No. 7**

**1<sup>st</sup>  
Movement**

**: Introduction and Basic Strategy (10 min)**

# Cracow Symphony No. 7

**1<sup>st</sup>  
Movement**

: **Introduction and Basic Strategy (10 min)**

**2<sup>nd</sup>  
Movement**

: **Expectations and first Messages from  
New Animalcula (15 min)**

# Cracow Symphony No. 7

**1<sup>st</sup>  
Movement**

: **Introduction and Basic Strategy (10 min)**

**2<sup>nd</sup>  
Movement**

: **Expectations and first Messages from  
New Animalcula (15 min)**

**3<sup>rd</sup>  
Movement**

: **New Animalcula Fairytales (10 min)**

# Cracow Symphony No. 7

**1<sup>st</sup>  
Movement**

: **Introduction and Basic Strategy (10 min)**

**2<sup>nd</sup>  
Movement**

: **Expectations and first Messages from  
New Animalcula (15 min)**

**3<sup>rd</sup>  
Movement**

: **New Animalcula Fairytales (10 min)**

**4<sup>th</sup>  
Movement**

: **Finale: Vivace ! (2 min)**

(hep-ph/0910.1032): “Flavour Theory : 2009”

(hep-ph/1012.1447 ): “MFV and Beyond”


# **1st Movement**

## **Introduction and Basic Strategy**

**CKM**

**(Nobel Prize 2008)**


**Dirac Medal  
(2010)**




**N. Cabibbo  
(1935-2010)**



**M. Kobayashi**



**T. Maskawa**




**EPS**


(High Energy Prize 2011)



**Sheldon Glashow**



**John Iliopoulos**




**Luciano Maiani**

**GIM**

# Asymptotic Freedom

(Nobel Prize 2004)  
(EPS High Energy  
Prize 2003)



**David Gross**




**Frank Wilczek**



**David Politzer**

# Asymptotic Freedom in QCD



Bethke  
hep-ex/0908.1135

Gross  
Politzer  
Wilczek  
(1973)

Nobel Prize  
2004

# Hierarchical Structure of the CKM Matrix

$$\begin{pmatrix} 0.97 & s_{12} & s_{13} e^{-iy} \\ -s_{12} & 0.97 & s_{23} \\ s_{12}s_{23} - s_{13}e^{iy} & -s_{23} & 1 \end{pmatrix}$$

$$s_{13} \ll s_{23} \ll s_{12}$$

$$(4 \cdot 10^{-3}) \quad (4 \cdot 10^{-2}) \quad (0.2)$$



## GIM Structure of FCNC's

**Large  $\mathcal{CP}$  effects in  $B_d$**

**Small  $\mathcal{CP}$  effects in  $B_s$**

**Tiny  $\mathcal{CP}$  effects in  $K_L$**

$$A_{CP}(B_d \rightarrow \psi K_s) \approx 0(1)$$

$$S_{\psi K_s} \approx \frac{2}{3}$$

$$A_{CP}(B_s \rightarrow \psi \phi) \approx 0(10^{-2})$$

$$S_{\psi \phi} \approx \frac{1}{25}$$

$$\varepsilon \approx 0(10^{-3}) \quad \varepsilon' \approx 0(10^{-6})$$

$$Br(K_L \rightarrow \pi^0 \bar{v} v) \approx 0(10^{-11})$$

## PMNS: Negligible LFV

(tiny  $v$  masses)


# Crucial Question

**What is the Origin of  
Particle Masses and the Reason  
for their Hierarchy and  
Hierarchy of their  
Flavour-Changing Interactions ?**

**Which Dynamics could be responsible for the observed  
structure of Electroweak Symmetry Breaking and of  
Patterns seen in Flavour Physics ?**

- 1.** Could it be an elementary SM Higgs system with all problems of instability under radiative corrections (hierarchy problems) ?  
Crucial questions in Particle Physics
- 2.** Could it be a new strong dynamics with a composite Higgs or without Higgs at all ?
- 3.** Could this dynamics help us understanding matter-antimatter asymmetry and the amount of dark matter in the universe ?
- 4.** Would these dynamics explain anomalies in flavour physics ?

# Indirect Search: Precision Measurements of Decays of Mesons and Leptons




$$B^+ \rightarrow \tau^+ \nu_\tau$$

Standard Model

$$\text{Br}(B^+ \rightarrow \tau^+ \nu_\tau)_{\text{SM}} = \left| A \frac{g_w^2}{M_w^2} \right|^2$$

$m_B \approx 5 \text{ GeV}$

A, B – parameters of a given theory



Contribution  
of a new  
charged Heavy  
Particle

$$\begin{aligned} \text{Br}(B^+ \rightarrow \tau^+ \nu_\tau) \\ = \left| A \frac{g_w^2}{M_w^2} + B \frac{g_H^2}{M_H^2} \right|^2 \end{aligned}$$

$$\Delta = \text{Br}(B^+ \rightarrow \tau^+ \nu_\tau) - \text{Br}(B^+ \rightarrow \tau^+ \nu_\tau)_{\text{SM}} \neq 0$$

Signal of a  
new particle

## In Order to identify New Animalcula through Flavour Physics

We need

- 1.** Many precision measurements of many observables and precise theory.
- 2.** Study Patterns on Flavour Violation in various New Physics models (correlations between many flavour observables).

...and

3.

### **Correlations between low energy flavour observables and Collider Physics (LHC, Tevatron)**

**Here top-down approach more  
powerful in flavour physics**

# Basic Questions for Flavour Physics

New Flavour  
violating  
CPV phases?

Flavour Conserving  
CPV phases?

Non-MFV  
Interactions?

(Non-CKM)

Right-Handed  
Charged  
Currents?


Scalars  $H^0, H^\pm$   
and related  
FCNC's?

New Fermions?  
New Gauge  
Bosons?




How to explain dynamically 22 free  
Parameters in the Flavour Sector ?

# Most popular BSM Directions




New gauge bosons, fermions, scalars in loops  
and even trees with often non-CKM interactions.

# Basic Diagrams in FCNC Processes



## Patterns of Flavour Violation around the Flavour Clock



# Superstars of 2012 – 2018 (Flavour Physics)

$S_{\psi\varphi}$   
 $\mathcal{CP}$  in  $B_s^0 - \bar{B}_s^0$

$$(B_s \rightarrow \varphi\varphi)$$

$B_s \rightarrow \mu^+ \mu^-$   
 $(B_d \rightarrow \mu^+ \mu^-)$

$$\begin{aligned} B &\rightarrow K^* \mu^+ \mu^- \\ B &\rightarrow X_s \mu^+ \mu^- \end{aligned}$$

$K^+ \rightarrow \pi^+ \nu\bar{\nu}$   
 $(K_L \rightarrow \pi^0 \nu\bar{\nu})$

$$\begin{aligned} B &\rightarrow K^* \nu\bar{\nu} \\ B &\rightarrow X_s \nu\bar{\nu} \end{aligned}$$

$\gamma, V_{ub}$   
from Tree  
Level  
Decays

$$\begin{aligned} \mu &\rightarrow e\gamma \\ \tau &\rightarrow \mu\gamma \\ \tau &\rightarrow e\gamma \\ \mu &\rightarrow 3e \\ \tau &\rightarrow 3 \text{ leptons} \end{aligned}$$

$$B^+ \rightarrow \tau^+ \nu_\tau$$

EDM's  
 $(g-2)_\mu$

$$\varepsilon'/\varepsilon$$

(Lattice)


# **2nd Movement**

**Expectations and First Messages  
from New Animalcula**

# First Evidence for CP Violation in Charm

Difference  
in time-  
integrated  
CP asymmetries

$$\Delta A_{CP} \equiv A_{CP}(K^-K^+) - A_{CP}(\pi^+\pi^-)$$



$$\Delta A_{CP} \equiv -0.82 \pm 0.21(\text{stat}) \pm 0.11(\text{sys})\%$$

Significance  $3.5\sigma$ ; Sensitive mainly to direct CPV

**VERY  
PRELIMINARY**

Central value larger than SM expectation  
but theoretical uncertainties in direct CPV  
are substantial.

From  
Mat Charles (Oxford)  
LHCb-CONF-2011-061

# Departures from Standard Model Expectations

<b>CP</b>	$K^0 - \bar{K}^0$	$(\varepsilon_K)$	$\frac{ \varepsilon_K _{SM}}{ \varepsilon_K _{exp}} \approx 0.80 \pm 0.10$	(AJB, Guadagnoli) (Brod, Gorbahn)
	$B_d^0 - \bar{B}_d^0$	$(S_{\psi K_s})$	$(S_{\psi K_s}) \approx 0.82 \pm 0.04$ (SM)    (UTfit)	
	$B_s^0 - \bar{B}_s^0$	$(S_{\psi\varphi})$	$(S_{\psi\varphi})_{exp} \approx 0.678 \pm 0.022$ (exp)	
	$\frac{\text{Br}(B^+ \rightarrow \tau^+ \nu)_{exp}}{\text{Br}(B^+ \rightarrow \tau^+ \nu)_{SM}} \approx 2.2 \pm 0.5$	$\frac{(S_{\psi\varphi})_{exp}}{(S_{\psi\varphi})_{SM}} \approx 10 - 20$	(CDF, DØ, Lenz+Nierste)	Spring 2011
		 <b>0.04</b>	$(S_{\psi\varphi})_{exp} \approx 0.8^{+0.1}_{-0.2}$	
	$ V_{ub}  = \begin{cases} 4.4 \cdot 10^{-3} & \text{Inclusive Decays } (B \rightarrow X_u l \bar{\nu}) \\ 3.4 \cdot 10^{-3} & \text{Exclusive Decays } (B \rightarrow \rho l \bar{\nu}) \\ & \text{and SM-CKM fit} \end{cases}$			(Right-handed currents? Crivellin; Mannel et al. AJB, Gemmeler, Isidori)

# News about New Physics from Summer Conferences

DØ, CDF, LHCb

$$-0.1 \leq S_{\psi\varphi} \leq 0.4 \quad *)$$



\*) Altmannshofer + Carena  
**1110.0843**

Can  $|V_{ub}|_{\text{excl}} \neq |V_{ub}|_{\text{incl}}$  be explained through right-handed currents?

Crivellin; Chen + Nam; Feger, Mannel et al.; AJB, Gemmeler, Isidori

$$|V_{ub}|_{\text{excl}} = 3.38 \ (36) \cdot 10^{-3}$$

$$|V_{ub}|_{\text{inc}} = 4.27 \ (38) \cdot 10^{-3}$$

$$\varepsilon \approx \frac{v_L}{v_R}$$

$$|V_{ub}|_{\text{excl}} = |V_{ub}^L + a\varepsilon^2 V_{ub}^R|$$

$$|V_{ub}|_{\text{inc}} \approx |V_{ub}^L|$$

Generally: in principle yes

But a very detailed analysis of  $SU(2)_L \otimes SU(2)_R \otimes U(1)_{B-L}$  with  $g_L \neq g_R$ ;  $V_L \neq V_R$  (mixing) including FCNC constraints + EWP constraints shows that in this concrete model the effect of RH currents too small !!

Blanke  
AJB  
Gemmeler  
Heidsieck  
1111.5014

## Two Scenarios for $|V_{ub}|$

(Taking into account  $\Delta M_s, \Delta M_d \leftarrow B_{d,s}^0 - \bar{B}_{d,s}^0$  Mixing)

$$\left\{ |V_{ub}| \approx 4.3 \cdot 10^{-3} \right\} \Rightarrow \left\{ \frac{\left( S_{\psi K_s} \right)_{SM}}{\left( S_{\psi K_s} \right)_{exp}} \right\} \approx 1.2 \quad \frac{|\varepsilon_K|_{SM}}{|\varepsilon_K|_{exp}} \approx 1.0$$
$$\left\{ |V_{ub}| \approx 3.4 \cdot 10^{-3} \right\} \Rightarrow \left\{ \frac{\left( S_{\psi K_s} \right)_{SM}}{\left( S_{\psi K_s} \right)_{exp}} \right\} \approx 1.0 \quad \frac{|\varepsilon_K|_{SM}}{|\varepsilon_K|_{exp}} \approx 0.8$$

New Physics  
in  $B_d^0 - \bar{B}_d^0$   
required

New Physics  
in  $\varepsilon_K$  required




Unfortunately to resolve  
this issue we have to wait  
for Belle II, Super-B and  
smarter Theorists

The size of CP Violation depends  
on the size of CKM elements:  
here  $|V_{ub}|$

# Possible Simplest Solutions

A

New negative CP phase  $\varphi_{\text{new}}$  in  $B_d^0 - \bar{B}_d^0$  Mixing  
 →  $|V_{ub}|$  from inclusive decays is correct



$$(S_{\psi K_s})_{\text{SM}} = \sin 2\beta \rightarrow S_{\psi K_s} = \sin(2\beta - \varphi_{\text{new}})$$

0.82

0.68

for  $\varphi_{\text{new}} = 10^\circ$

Soni, Lunghi

AJB, Guadagnoli  
 UTfitters  
 Lenz, Nierste +  
 CKMfitters  
 Laiho, Lunghi,  
 van der Water  
 Fleischer et al  
 Blanke et al  
 Branco et al

....



$\varepsilon_K$  and  $\text{Br}(B^+ \rightarrow \tau^+ \nu)$  much closer to experiment

B

Dynamical Model : **Non-Supersymmetric Two-Higgs Doublet Model with Flavour Blind Phases** (AJB, Carlucci, Gori, Isidori AJB, Isidori, Paradisi) **2HDM<sub>MFV</sub>**

Correlated Implications:



Enhanced  $S_{\psi\varphi}$ ,  $\text{Br}(B_s \rightarrow \mu^+ \mu^-)$ ,  $\text{Br}(B_d \rightarrow \mu^+ \mu^-)$ , EDM's

(non-SUSY)

## General 2HDM with MFV and Flavour Blind CPV Phases (in Yukawa Couplings)

(1005.5310)

(AJB, Carlucci, Gori, Isidori)

Provides correct pattern

$$\varepsilon_K : \quad \text{Diagram} \quad \approx \left[ \frac{m_d m_s}{M_H^2} \right] m_t^4 (\tan \beta)^2 (V_{ts}^* V_{td})^2 \quad (\text{tiny})$$

$$S_{\psi K_s} : \quad \text{Diagram} \quad \approx \left[ \frac{m_b m_d}{M_H^2} \right] m_t^4 (\tan \beta)^2 (V_{tb}^* V_{td})^2 e^{i\phi_{\text{new}}}$$

$$S_{\psi \varphi} : \quad \text{Diagram} \quad \approx \left[ \frac{m_b m_s}{M_H^2} \right] m_t^4 (\tan \beta)^2 (V_{tb}^* V_{ts})^2 e^{i\phi_{\text{new}}}$$

$$S_{\psi K_s} = \sin(2\beta - \theta_d^H) \quad S_{\psi \varphi} \approx \sin(\theta_s^H)$$

$$\sin 2\beta > S_{\psi K_s}$$

( $|\varepsilon_K|$  enhanced)

$$\frac{\theta_d^H}{\theta_s^H} \approx \frac{m_d}{m_s} \approx \frac{1}{17}$$

$$\begin{aligned} \tan \beta &\approx 10 - 20 \\ M_H &\approx 250 \text{ GeV} \end{aligned}$$

Large  
RG QCD  
effects  
 $Q_{LR}$

$|\epsilon_K|$  vs  $S_{\psi\phi}$  and  $S_{\psi K_s}$  vs  $S_{\psi\phi}$   
 in a General 2HDM with MFV and Flavour Blind CPV


(AJB, Carlucci, Gori, Isidori)

Correct pattern of NP effects

2HDM<sub>MFV</sub>

Correlation between various CP Effects


(But the effects appear a bit too weak)



1005.5310

# More on 2HDM with MFV and Flavour Blind Phases

## Correlation between CP Effects



AJB, Isidori, Paradisi 1007.5291

$$S_{\psi K_s} = \sin(2\beta - \theta_d^H) \quad S_{\psi \phi} \approx \sin(\theta_s^H)$$

$L_{\text{Yukawa}}$ :

$$\frac{\theta_d^H}{\theta_s^H} \approx \frac{m_d}{m_s} \approx \frac{1}{17}$$

BCGI

$L_{\text{Higgs}}$ :  
(potential)

$$\frac{\theta_d^H}{\theta_s^H} = 1$$

Kagan, Perez, Volansky, Zupan  
Paradisi, Straub  
Dobrescu, Fox, Martin  
Blum, Hochberg, Nir  
Ligeti, Papucci, Perez, Zupan

## Insight after Summer Conferences

$$\{-0.1 \leq S_{\psi\phi} \leq 0.4\} \Rightarrow \left\{ \begin{array}{l} \text{Phases in} \\ \text{Higgs Potential} \\ \text{favoured} \end{array} \right\}$$

LHCb, CDF, DØ

See also: Altmannshofer + Carena  
1110.0843 (MFV-MSSM  
+ higher-dimension  
operators)

**But  $|V_{ub}|$  could turn out to be small !**

$$|V_{ub}| \simeq |V_{ub}|_{\text{exl}} \approx 3.4 \cdot 10^{-3}$$

AJB, Guadagnoli  
(2008)

Then  $(S_{\psi K_s})_{\text{SM}} \simeq (S_{\psi K_s})_{\text{exp}}$

**Solution**

**C**

**But  $(\varepsilon_K)_{\text{SM}} \simeq 0.8(\varepsilon_K)_{\text{exp}}$**





**Need new contributions to  $\varepsilon_K$   
without new phases in  $B_d^0 - \bar{B}_d^0$**

**mixing**

AJB, Carlucci, Merlo, Stamou  
hep-ph 1112.4477 (2011)

**Gauged Flavour  $SU(3)^3$  Model of Grinstein et al  
provides an example**

# Correlations: $\varepsilon_K$ & $\Delta M_{B_d}$ and $\varepsilon_K$ & $\Delta M_{B_s}$ in Gauged SU(3)<sup>3</sup> Flavour Models



Solution to  $\varepsilon_K$  problem destroys the agreement  
with the data for  $\Delta M_{B_d}$  and  $\Delta M_{B_s}$

AJB, Carlucci, Merlo, Stamou  
hep-ph 1112.4477 (2011)



$$B_s \rightarrow \mu^+ \mu^- \text{ and } B_d \rightarrow \mu^+ \mu^-$$

Z-Penguin (SM  
+ Boxes CMFV)

SM

$$\text{Br}(B_s \rightarrow \mu^+ \mu^-) = (3.2 \pm 0.2) \cdot 10^{-9}$$

$$\text{Br}(B_d \rightarrow \mu^+ \mu^-) = (1.0 \pm 0.1) \cdot 10^{-10}$$

Error  
dominated  
by  $\hat{B}_{d,s}$

AJB (03)

CMFV  
“Golden  
Relation”

$$\frac{\text{Br}(B_s \rightarrow \mu^+ \mu^-)}{\text{Br}(B_d \rightarrow \mu^+ \mu^-)} = \frac{\hat{B}_d}{\hat{B}_s} \frac{\tau(B_s)}{\tau(B_d)} \frac{\Delta M_s}{\Delta M_d}$$

$(\Delta B = 1)$        $(0.95 \pm 0.03)$        $(\Delta B = 2)$   
Lattice

Valid in  
all CMFV  
models

Can be  
strongly  
violated  
in SUSY,  
LHT, RS,  
4G


95% CL

$$\text{Br}(B_s \rightarrow \mu^+ \mu^-) = \begin{cases} (18^{+11}_{-9}) \cdot 10^{-9} & (\text{CDF}) \\ < 11 \cdot 10^{-9} & (\text{LHC}) \end{cases}$$

$$\text{Br}(B_d \rightarrow \mu^+ \mu^-) \leq 4 \cdot 10^{-9} \quad (\text{LHC})$$

$$(\text{CDF})$$

# $B_s \rightarrow \mu^+ \mu^-$ Beyond the Standard Model



Other Z-Penguins  
and Boxes

SM:  $(3.2 \pm 0.2) \cdot 10^{-9}$

Model Independent  
Limit (95% C.L.)

$$\text{Br}(B_s \rightarrow \mu^+ \mu^-) < 5.6 \cdot 10^{-9}$$

Altmannshofer, Paradisi,  
Straub 1111.1257

$$\frac{(\tan \beta)^6}{M_H^4}$$

in SUSY

$$\text{Br}(B_s \rightarrow \mu^+ \mu^-) < 11 \cdot 10^{-9}$$



In the case of

$$\text{Br}(B_s \rightarrow \mu^+ \mu^-) > 6 \cdot 10^{-9}$$

distinction between  $Z, Z'$  and  $H^0$   
possible

# More on 2HDM with MFV and Flavour Blind Phases

2HDM<sub>MFV</sub>




AJB, Isidori, Paradisi 1007.5291

# **3rd Movement**

**New Animalcula  
Fairytales**

# Most popular BSM Directions



New gauge bosons, fermions, scalars in loops  
and even trees with often non-CKM interactions.

# Models with non-MFV Interactions facing Large $S_{\psi\phi}$

## Model Expectations

$$S_{\psi\phi} \leq \begin{cases} 0.80 & (4G) \text{ (Fourth Generation)} (t') \text{ (Soni, Hou, Munich, Lenz)} \\ 0.75 & (AC) \text{ (abelian flavour, SUSY)} \text{ (Higgs penguin)} \quad \text{ABGPS} \\ 0.50 & (\text{RVV}) \text{ (non - abelian flavour, SUSY)} \text{ (Higgs penguin)} \\ 0.75 & (\text{RS}) \quad (\text{Heavy KK Gauge Bosons}) \quad (\text{Duling et al (08)}) \\ 0.30 & (\text{LHT}) \text{ (Mirror Fermions at work)} \text{ (Tarantino et al (09))} \end{cases}$$

$$(S_{\psi\phi})_{\text{SM}} \approx 0.04$$

**ABGPS** = Altmannshofer, AJB, Gori, Paradisi, Straub  
**0909.1333**

# Implications of an Enhanced $S_{\psi\phi}$

- 1.** Enhanced  $\text{Br}(B_s \rightarrow \mu^+ \mu^-)$   
(SUSY flavour models, 2HDM<sub>MFV</sub>, 4G)
- 2.** Enhanced  $\text{Br}(B_d \rightarrow \mu^+ \mu^-)$   
(2HDM<sub>MFV</sub>, also in some SUSY flavour models)
- 3.**  $\text{Br}(B_d \rightarrow \mu^+ \mu^-)$  forced to be SM - like in 4G
- 4.**  $K^+ \rightarrow \pi^+ v\bar{v}$  and  $K_L \rightarrow \pi^0 v\bar{v}$  forced to be SM - like  
(LHT, Randall - Sundrum)
- 5.** Automatic enhancements in SUSY - GUT models:  
 $\text{Br}(\mu \rightarrow e\gamma)$ ,  $\text{Br}(\tau \rightarrow \mu\gamma)$ ,  $(g-2)_\mu$ ,  $d_e$ ,  $d_n$

**ABGPS**  
(0909.1333)



**$\text{Br}(B_d \rightarrow \mu^+ \mu^-) \text{ vs } \text{Br}(B_s \rightarrow \mu^+ \mu^-)$**

**SUSY**

 = SM

**MFV**

AJB; Hurth, Isidori, Kamenik, Mescia



**RVV2** (RH currents)

**LH currents**

CDF, D0  
LHCb



$\text{Br}(B_s \rightarrow \mu^+ \mu^-) \text{vs } S_{\psi\phi}$

SUSY

ABGPS

(0909.1333)

$\Delta a_\mu \sim \Delta(g - 2)_\mu \text{ vs } S_{\psi\phi}$



$$K^+ \rightarrow \pi^+ \nu\bar{\nu} \text{ and } K_L \rightarrow \pi^0 \nu\bar{\nu} \text{ (Z°-penguins)}$$

(TH cleanest FCNC decays in Quark Sector)

Extensive  
TH efforts  
over  
20 years

: Buchalla, AJB; Misiak, Urban (NLO QCD)  
AJB, Gorbahn, Haisch, Nierste (NNLO QCD)  
Brod, Gorbahn, Stamou (QED, EW two loop)  
Isidori, Mescia, Smith (several LD analyses)  
Buchalla, Isidori (LD in  $K_L \rightarrow \pi^0 \nu\bar{\nu}$ )

$$\frac{\text{Br}(K^+ \rightarrow \pi^+ \nu\bar{\nu})}{\text{Br}(K_L \rightarrow \pi^0 \nu\bar{\nu})} = 3.2 \pm 0.2$$

**SM** :  $\text{Br}(K^+ \rightarrow \pi^+ \nu\bar{\nu}) = (8.4 \pm 0.7) \cdot 10^{-11}$

$$\text{Br}(K_L \rightarrow \pi^0 \nu\bar{\nu}) = (2.6 \pm 0.4) \cdot 10^{-11}$$

**Exp** :  $\text{Br}(K^+ \rightarrow \pi^+ \nu\bar{\nu}) = \left( 17^{+11}_{-10} \right) \cdot 10^{-11}$

$$\text{Br}(K_L \rightarrow \pi^0 \nu\bar{\nu}) \leq 6.8 \cdot 10^{-8}$$

(E787, E949      Brookhaven)

(E391a, KEK)

Future :

NA62  
Project X (FNAL)

Both very  
sensitive to  
New Physics

CP-conserving  
TH uncertainty 2-3%

J-PARC KOTO

CP-Violation in Decay  
TH uncertainty 1-2%

## Important Messages

**1.**

**Many Models (SUSY, 4G, LHT, RS)  
can still accommodate**

$$\text{Br}(K^+ \rightarrow \pi^+ \nu\bar{\nu}) \sim 3 \text{Br}(K^+ \rightarrow \pi^+ \nu\bar{\nu})_{\text{SM}}$$
$$\text{Br}(K_L \rightarrow \pi^0 \nu\bar{\nu}) \sim 10 \text{Br}(K_L \rightarrow \pi^0 \nu\bar{\nu})_{\text{SM}}$$

**2.**

**Even if no significant New Physics  
would be seen in B-decays  
large effects in  $K \rightarrow \pi \nu\bar{\nu}$  are possible.**

# Lepton Flavour Violation, $\Delta(g - 2)_\mu$ and EDM's

(MEGA)  $Br(\mu \rightarrow e\gamma) < 1.2 \cdot 10^{-11}$   $\rightarrow 10^{-13}$  (MEG) SM:  $10^{-54}$

$$(a_\mu)_{\text{SM}} < (a_\mu)_{\text{exp}} \quad (3.1\sigma)$$

$$a_\mu = \frac{1}{2} (\mathbf{g} - 2)_\mu$$

(Regan et al)  $d_e < 1.6 \cdot 10^{-27}$   $\rightarrow 10^{-31}$   $(d_e)_{\text{SM}} \approx 10^{-38}$

[e cm]

(Baker et al)  $d_n < 2.9 \cdot 10^{-26}$   $\rightarrow 10^{-28}$   $(d_n)_{\text{SM}} \approx 10^{-32}$

# Lepton Flavour Violation, $\Delta(g - 2)_\mu$ and EDM's

(MEGA)  $Br(\mu \rightarrow e\gamma) < 1.2 \cdot 10^{-11}$   $\rightarrow 10^{-13}$  (MEG) SM:  $10^{-54}$

$$(a_\mu)_{\text{SM}} < (a_\mu)_{\text{exp}} \quad (3.1\sigma)$$

$$a_\mu = \frac{1}{2} (\mathbf{g} - 2)_\mu$$

(Regan et al)  $d_e < 1.6 \cdot 10^{-27}$   $\rightarrow 10^{-31}$   $(d_e)_{\text{SM}} \approx 10^{-38}$



[e cm]

(Baker et al)  $d_n < 2.9 \cdot 10^{-26}$   $\rightarrow 10^{-28}$   $(d_n)_{\text{SM}} \approx 10^{-32}$





MEG:  $Br(\mu \rightarrow e\gamma) \leq 2 \cdot 10^{-12}$

# Correlations in the SU(3) Flavour SUSY Model (RVV)



MEG

# Correlations within SUSY-SU(5)-GUT with RH Neutrinos



AJB, Nagai, Paradisi, 1011.1993

# DNA Tests of Flavour Models

$O_i$  : *Observables*

$M_i$  : *Models beyond SM*

	$M_1$	$M_2$	$M_3$	$M_4$	$M_5$
$O_1$	★★★	★	★	★	★★
$O_2$	★	★★	★★★	★★	★
$O_3$	★★	★★★	★★	★	★
$O_4$	★★★	★★	★	★★★	★★
$O_5$	★	★★★	★	★★	★★★



Very large New Physics effect



Moderate New Physics effect



Very small New Physics effect

	AC	RVV2	AKM	$\delta LL$	FBMSSM	LHT	RS	4G
$D^0 - \bar{D}^0$	★★★	★	★	★	★	★★★	?	★★
$\epsilon_K$	★	★★★	★★★	★	★	★★	★★★	★★
$S_{\psi\phi}$	★★★	★★★	★★★	★	★	★★★	★★★	★★★
$S_{\phi K_S}$	★★★	★★	★	★★★	★★★	★	?	★★
$A_{CP}(B \rightarrow X_s \gamma)$	★	★	★	★★★	★★★	★	?	★
$A_{7,8}(B \rightarrow K^* \mu^+ \mu^-)$	★	★	★	★★★	★★★	★★	?	★★
$A_9(B \rightarrow K^* \mu^+ \mu^-)$	★	★	★	★	★	★	?	★★
$B \rightarrow K^{(*)} \nu \bar{\nu}$	★	★	★	★	★	★	★	★
$B_s \rightarrow \mu^+ \mu^-$	★★★	★★★	★★★	★★★	★★★	★	★	★★★
$K^+ \rightarrow \pi^+ \nu \bar{\nu}$	★	★	★	★	★	★★★	★★★	★★★
$K_L \rightarrow \pi^0 \nu \bar{\nu}$	★	★	★	★	★	★★★	★★★	★★★
$\mu \rightarrow e \gamma$	★★★	★★★	★★★	★★★	★★★	★★★	★★★	★★★
$\tau \rightarrow \mu \gamma$	★★★	★★★	★	★★★	★★★	★★★	★★★	★★★
$\mu + N \rightarrow e + N$	★★★	★★★	★★★	★★★	★★★	★★★	★★★	★★★
$d_n$	★★★	★★★	★★★	★★	★★★	★	★★★	★
$d_e$	★★★	★★★	★★	★	★★★	★	★★★	★
$(g-2)_\mu$	★★★	★★★	★★	★★★	★★★	★	?	★




# Cracow 2020 Vision

	NEW SM
$D^0 - \bar{D}^0$	★★
$\epsilon_K$	★★
$S_{\psi\phi}$	★★★
$S_{\phi K_S}$	★★
$A_{\text{CP}}(B \rightarrow X_s \gamma)$	★
$A_{7,8}(B \rightarrow K^* \mu^+ \mu^-)$	★★
$A_9(B \rightarrow K^* \mu^+ \mu^-)$	★
$B \rightarrow K^{(*)} \nu \bar{\nu}$	★★★
$B_s \rightarrow \mu^+ \mu^-$	★★★
$K^+ \rightarrow \pi^+ \nu \bar{\nu}$	★★
$K_L \rightarrow \pi^0 \nu \bar{\nu}$	★★★
$\mu \rightarrow e \gamma$	★★★
$\tau \rightarrow \mu \gamma$	★★★
$\mu + N \rightarrow e + N$	★★★
$d_n$	★★★
$d_e$	★★★
$(g-2)_\mu$	★★



# **4th Movement**

**Finale: Vivace !**



# Superstars of 2012 – 2018 (Flavour Physics)

$S_{\psi\phi}$   
 $\mathcal{CP}$  in  $B_s^0 - \bar{B}_s^0$

$$(B_s \rightarrow \phi\phi)$$

$B_s \rightarrow \mu^+ \mu^-$   
 $(B_d \rightarrow \mu^+ \mu^-)$

$$\begin{aligned} B &\rightarrow K^* \mu^+ \mu^- \\ B &\rightarrow X_s \mu^+ \mu^- \end{aligned}$$

$K^+ \rightarrow \pi^+ \nu\bar{\nu}$   
 $(K_L \rightarrow \pi^0 \nu\bar{\nu})$

$$\begin{aligned} B &\rightarrow K^* \nu\bar{\nu} \\ B &\rightarrow X_s \nu\bar{\nu} \end{aligned}$$

$\gamma, V_{ub}$   
from Tree  
Level  
Decays

$$\begin{aligned} \mu &\rightarrow e\gamma \\ \tau &\rightarrow \mu\gamma \\ \tau &\rightarrow e\gamma \\ \mu &\rightarrow 3e \\ \tau &\rightarrow 3 \text{ leptons} \end{aligned}$$


$$B^+ \rightarrow \tau^+ \nu_\tau$$

EDM's  
 $(g-2)_\mu$

$$\varepsilon'/\varepsilon$$

(Lattice)

## Patterns of Flavour Violation around the Flavour Clock



# Many Thanks to my Collaborators

SUSY



W. Altmannshofer



S. Gori



P. Paradisi



D. Straub

LHT



M. Blanke



B. Duling



A. Poschenrieder



S. Recksiegel



C. Tarantino



S. Uhlig



A. Weiler

RS



M. Albrecht



M. Blanke



B. Duling



K. Gemmler



S. Gori



A. Weiler

# ALL WANTED !!

SUSY



W. Altmannshofer



S. Gori



P. Paradisi



D. Straub


LHT



M. Blanke



B. Duling



A. Poschenrieder



S. Recksiegel



C. Tarantino



S. Uhlig



A. Weiler

RS



M. Albrecht



M. Blanke



B. Duling



K. Gemmler



S. Gori



A. Weiler

**4 G**



**B. Duling**



**T. Heidsieck**



**C. Promberger**



**T. Feldmann**



**S. Recksiegel**

**2 HDM**



**M.V. Carlucci**



**S. Gori**



**G. Isidori**

**$\epsilon_K$**



**D. Guadagnoli**

**RH Currents**



**K. Gemmeler**



**G. Isidori**


# Most Recent Collaborators



J. Girrbach



C. Grojean



S. Pokorski



R. Ziegler

Theory of  
Fermion Masses



M.V. Carlucci



L. Merlo



E. Stamou

Gauge  
Flavour  
Models



P. Paradisi



M. Nagai



M. Blanke

Cracow0112



K. Gemmeler



T. Heidsieck

LR  
Models

2HDM $\overline{\text{MFV}}$

# **Should we be frustrated after Summer Conferences ?**

**Should we be frustrated  
after Summer Conferences ?**

**No, no, no !!!**

**Should we be frustrated  
after Summer Conferences ?**

**No, no, no !!!**

**Exciting Times are just  
ahead of us !!!**

# **New Animalcula in Sight !**

# **New Animalcula in Sight !**

**Hopefully we will soon know  
how they really look like !**

# **Thank You !!**

## **New Animalcula in Sight !**

**Hopefully we will soon know  
how they really look like !**



# Backup

Similar  
Result  
by Soni et al.

$\text{Br}(B_s \rightarrow \mu^+ \mu^-) \text{ vs } S_{\psi\phi}$

4G

BDFHPR  
(1002.2126)



No Impact  
on  $\Delta a_\mu$

4G has hard time to describe simultaneously  $\epsilon'/\epsilon$  and  $S_{\psi\phi} > 0.2$  if  $B_{6,8}$  within 20% from large N values